

Length-weight Relationships of Nigerian Freshwater Fishes

Richard P. King

Abstract

Length-weight relationships of 73 fish populations, covering 20 families, 28 genera and 40 species inhabiting freshwater ecosystems in Nigeria, were estimated (73 cases) or assembled from the literature (20 cases), and tested for difference between ecosystem types. There were no significant differences in the exponent of these relationships between lotic and lentic systems.

Introduction

Length-weight relationships (LWR) of fish are important in fisheries biology because they allow inter alia:

- (i) estimation of average weight of the fish of a given length group (Beyer 1987);
- (ii) conversion of length-growth equations to weight-growth equivalents (i.e., length-at-age to weight-at-age) in yield-per-recruit and related models;
- (iii) interspecific and interpopulational morphometric comparison of fish species; and
- (iv) assessing the relative well-being of fish populations (Bolger and Connolly 1989).

Only limited species-specific LWR data are available for the freshwater fish resource of Nigeria, West Africa, and the present contribution is aimed at partly compensating for this.

Materials and Methods

Fishes were sampled from different freshwater bodies (both lotic and lentic systems) in Nigeria from 1984 to 1994, using a variety of methods/gears, including gill-nets, dip-nets, hooks and traditional valved basket traps. They were identified, measured (cm, total or standard length) and weighted (g, total fresh weight). Nomenclature of the fish taxa conforms to Lévêque et al. (1992).

For each species, the parameters a (proportionality constant) and b (exponent) of the LWR of the form:

$$W = a \cdot L^b \quad \dots 1)$$

were estimated through base-10 logarithm transformation of L-W data pairs and ordinary least-squares linear regression (i.e., log transformed versions of equation 1) viz:

$$\log W = \log a + b \log L \quad \dots 2)$$

Whenever possible, LWR were determined separately, and for both sexes (including immature fishes). These different estimates are being treated here as separate "populations". Additional LWR parameters were obtained from the literature; in some cases here, important information was missing, e.g., sample sizes, correlation coefficients and/or the range of sizes considered (though in some cases, the size ranges could be read off graphs).

Results and Discussion

The LW data of 73 inland water fish populations, from 20 families, 28 genera and 40 species were analyzed.

Two species (*Mondactylus sebae* and *Bostrycus africana*) of brackishwater origin but which consistently maintained populations in fresh waters were included in the analyses.

Table 1. Length-weight relationships and related statistics of 73 populations of fish occurring in the inland waters of Nigeria.

| Family/Species | Sex | Length type | Length (cm) | | a | b | N | r | Area | Reference |
|-----------------------------------|-----|----------------|-------------|-------|---------|-------|-----|--------|-----------------|----------------------------|
| | | | min | max | | | | | | |
| Polypetridae | | | | | | | | | | |
| <i>Erpetoichthys calabaricus</i> | | T | 22.3 | 34.2 | 0.0033 | 2.749 | 152 | 0.977 | Mfangmfang pond | this study |
| <i>Erpetoichthys calabaricus</i> | | T | 21.6 | 41.4 | 0.0010 | 3.103 | 180 | 0.969 | Cross River | this study |
| <i>Erpetoichthys calabaricus</i> | | T | 20.1 | 33.6 | 0.0050 | 2.581 | 146 | 0.933 | Qua Iboe River | this study |
| Notopteridae | | | | | | | | | | |
| <i>Papyrocranus afer</i> | | S | 10.3 | 39.8 | 0.00004 | 2.663 | 162 | 0.979 | Ikpa River | this study |
| Momyridae | | | | | | | | | | |
| <i>Brienomyrus brachyistius</i> | | T | 3.1 | 16.0 | 0.1821 | 2.538 | 50 | 0.972 | Ikpa River | King (1989) |
| <i>Isichthys henryi</i> | | T | 20.0 | 31.9 | 0.0267 | 2.252 | 148 | 0.997 | Imo River | this study |
| <i>Petrocephalus ansorgii</i> | M | T | 10.9 | 15.0 | 0.0075 | 3.125 | 142 | 0.942 | Ikpa River | this study |
| <i>Petrocephalus ansorgii</i> | F | T | 10.9 | 15.2 | 0.0629 | 2.299 | 160 | 0.883 | Ikpa River | this study |
| Gymnarchidae | | | | | | | | | | |
| <i>Gymnarchus niloticus</i> | | T | 50.0 | 150.0 | 0.0039 | 2.978 | 58 | 0.985 | Lake Chad | Sagua (1986) |
| Characidae | | | | | | | | | | |
| <i>Brycinus longipinnis</i> | | T | 6.5 | 11.0 | 0.0129 | 3.109 | 60 | 0.932 | Ikpa River | Oni et al. (1983) |
| <i>Brycinus longipinnis</i> | | T | 7.5 | 11.6 | 0.0025 | 3.733 | 42 | 0.947 | Qua Iboe River | Oni et al. (1983) |
| <i>Brycinus macrolepidotus</i> | | T | 12.6 | 24.7 | 0.0137 | 2.906 | 33 | 0.999 | Ikpa River | Oni et al. (1983) |
| <i>Brycinus nurse</i> | | S | - | - | 0.0390 | 2.668 | 10 | 0.998 | River Galma | Oni et al. (1983) |
| <i>Brycinus imberi</i> | M | T | 18.9 | 32.8 | 0.0315 | 2.688 | 148 | 0.971 | Cross River | Oni et al. (1983) |
| <i>Brycinus imberi</i> | F | T | 17.1 | 22.0 | 0.0093 | 3.143 | 135 | 0.953 | Cross River | Oni et al. (1983) |
| <i>Brycinus imberi</i> | | T | 17.1 | 32.8 | 0.0549 | 2.523 | 283 | 0.980 | Cross River | Oni et al. (1983) |
| <i>Brycinus imberi</i> | | T | 10.5 | 15.7 | 0.0260 | 2.722 | 136 | 0.954 | Ikpa River | Oni et al. (1983) |
| Hepsetidae | | | | | | | | | | |
| <i>Hepsetus odoe</i> | | T | 9.2 | 17.1 | 0.0069 | 2.986 | 47 | 0.964 | Adadama Lake | this study |
| <i>Hepsetus odoe</i> | | T | 13.0 | 35.5 | 0.0022 | 3.376 | 60 | 0.9950 | Ikpa River | this study |
| Distichodontidae | | | | | | | | | | |
| <i>Nannaethiops unitaeniatus</i> | | T | 3.2 | 5.8 | 0.0653 | 2.158 | 82 | 0.980 | Mfangmfang pond | this study |
| Cyprinidae | | | | | | | | | | |
| <i>Barbus callipterus</i> | | T | 4.9 | 7.0 | 0.0103 | 2.845 | 180 | 0.936 | Qua Iboe River | this study |
| <i>Barbus callipterus</i> | | T | 4.8 | 7.4 | 0.0254 | 2.603 | 96 | 0.864 | Mgangmfang pond | this study |
| Bagridae | | | | | | | | | | |
| <i>Parauchenoglanis akiri</i> | | T | 6.5 | 10.0 | 0.0125 | 2.974 | 87 | 0.994 | Qua Iboe River | this study |
| <i>Parauchenoglanis akiri</i> | | T | 7.2 | 16.7 | 0.0197 | 2.841 | 111 | 0.989 | Ikpa River | this study |
| <i>Parauchenoglanis fasciatus</i> | | T | 4.8 | 13.5 | 0.0312 | 2.627 | 147 | 0.939 | Ikpa River | this study |
| <i>Parauchenoglanis fasciatus</i> | | T | 12.4 | 18.0 | 0.0052 | 3.245 | 170 | 0.949 | Zaria | this study |
| <i>Chrysichthys auratus</i> | | T | 9.0 | 24.0 | 0.0160 | 2.880 | 70 | 0.974 | Zaria | Nwadiaro and Okorie (1985) |
| <i>Chrysichthys walkeri</i> | | T | 7.0 | 24.0 | 0.0074 | 3.114 | - | - | Zaria | Ikusemiju (1976) |
| Clariidae | | | | | | | | | | |
| <i>Clarias macromystax</i> | | T | 6.5 | 10.0 | 0.0125 | 2.974 | 87 | 0.994 | Qua Iboe River | this study |
| <i>Clarias buthupogon</i> | | T | 7.2 | 16.7 | 0.0197 | 2.841 | 111 | 0.989 | Ikpa River | this study |
| <i>Clarias buthupogon</i> | | T | 4.8 | 13.5 | 0.0312 | 2.627 | 147 | 0.939 | Ikpa River | this study |
| <i>Clarias gariepinus</i> | M | S | 12.4 | 18.0 | 0.0052 | 3.245 | 170 | 0.949 | Cross River | Olatunde (1985) |
| <i>Clarias gariepinus</i> | F | S | 9.0 | 24.0 | 0.0160 | 2.880 | 70 | 0.974 | Oguta Lake | Olatunde (1985) |
| <i>Clarias gariepinus</i> | | S | 7.0 | 24.0 | 0.0074 | 3.114 | - | - | Lekki Lagoon | Olatunde (1985) |

Continued...

Table 1. Continued.

| Family/Species | Sex | Length type | Length (cm) min | max | a | b | N | r | Area | Reference |
|----------------------------------|-----|-------------|-----------------|------|--------|-------|-----|-------|-------------------|--------------------------|
| Malapteruridae | | | | | | | | | | |
| <i>Malapterurus electricus</i> | | T | 7.7 | 17.7 | 0.0018 | 2.893 | 222 | 0.992 | Qua Iboe River | this study |
| <i>Malapterurus electricus</i> | | T | 14.5 | 28.8 | 0.0080 | 3.160 | 124 | 0.985 | Cross River | this study |
| <i>Malapterurus electricus</i> | | T | 11.4 | 22.0 | 0.0108 | 3.069 | 150 | 0.983 | Imo River | this study |
| Mochokidae | | | | | | | | | | |
| <i>Synodontis schall</i> | | S | - | - | 0.1380 | 2.333 | 10 | 0.999 | River Galma | Oni et al. (1983) |
| <i>Synodontis schall</i> | M | S | 7.9 | 17.8 | 0.0162 | 3.170 | 111 | 0.990 | Zaria | Olatunde (1989) |
| <i>Synodontis schall</i> | F | S | 9.8 | 19.1 | 0.0200 | 3.110 | 150 | 0.830 | Zaria | Olatunde (1989) |
| Schilbeidae | | | | | | | | | | |
| <i>Eutropius niloticus</i> | M | S | 10.0 | 20.0 | 0.0102 | 3.020 | 109 | 0.980 | Lake Kainji | Olatunde (1979) |
| <i>Eutropius niloticus</i> | F | S | 7.0 | 22.4 | 0.0091 | 3.070 | 474 | 0.990 | Lake Kainji | Olatunde (1979) |
| <i>Schilbe mystus</i> | M | S | 6.8 | 15.0 | 0.0148 | 2.930 | 26 | 0.990 | Lake Kainji | Olatunde (1979) |
| <i>Schilbe mystus</i> | F | S | 9.0 | 20.0 | 0.0063 | 3.240 | 111 | 0.980 | Lake Kainji | Olatunde (1979) |
| <i>Schilbe mystus</i> | | T | 14.7 | 20.1 | 0.0284 | 2.506 | 70 | 0.944 | Ikpa River | this study |
| Aplocheilidae | | | | | | | | | | |
| <i>Epiplatys sexfasciatus</i> | M | T | - | - | 0.0290 | 2.730 | 130 | - | Adada River | Inyang and Anozie (1987) |
| <i>Epiplatys sexfasciatus</i> | F | T | - | - | 0.0060 | 3.520 | 129 | - | Adada River | Inyang and Anozie (1987) |
| <i>Epiplatys sexfasciatus</i> | | T | - | - | 0.0140 | 2.660 | 146 | - | Adada River | Inyang and Anozie (1987) |
| <i>Aphyosemion splendopleura</i> | | T | 3.3 | 4.8 | 0.0179 | 2.683 | 39 | 0.865 | Mfangmfang pond | this study |
| <i>Aphyosemion gardneri</i> | | T | 1.5 | 5.3 | 0.0141 | 2.649 | 311 | 0.864 | Mfangmfang pond | this study |
| <i>Aphyosemion arnoldi</i> | | T | 2.7 | 7.4 | 0.0239 | 2.672 | 72 | 0.983 | New Calabar River | this study |
| <i>Aphyosemion sjoestedti</i> | | T | 2.8 | 4.9 | 0.0252 | 2.435 | 70 | 0.942 | New Calabar River | this study |
| Channidae | | | | | | | | | | |
| <i>Parachanna obscura</i> | | T | 3.8 | 33.9 | 0.0098 | 2.904 | 192 | 0.990 | Cross River | this study |
| <i>Parachanna obscura</i> | | T | 9.5 | 56.0 | 0.0063 | 3.134 | 40 | 0.983 | Adadama Lake | this study |
| <i>Parachanna obscura</i> | | T | 10.9 | 32.3 | 0.0059 | 3.088 | 160 | 0.989 | Imo River | this study |
| <i>Parachanna africana</i> | | T | 12.0 | 22.2 | 0.0065 | 3.041 | 112 | 0.957 | Ikpa River | this study |
| Monodactylidae | | | | | | | | | | |
| <i>Monodactylus sebae</i> | | T | 9.3 | 17.5 | 0.0489 | 2.799 | 67 | 0.987 | Ikpa River | this study |
| Cichlidae | | | | | | | | | | |
| <i>Chromidotilapia guntheri</i> | | T | 9.0 | 16.5 | 0.0343 | 2.843 | 111 | 0.991 | Mfangmfang pond | this study |
| <i>Chromidotilapia guntheri</i> | | T | 8.3 | 23.5 | 0.0142 | 3.179 | 128 | 0.910 | Qua Iboe River | this study |
| <i>Thysochromis ansorgii</i> | | T | 6.0 | 12.5 | 0.0273 | 2.845 | 118 | 0.901 | Mfangmfang pond | this study |
| <i>Thysochromis ansorgii</i> | | T | 4.5 | 14.4 | 0.0112 | 3.173 | 156 | 0.988 | Cross River | this study |
| <i>Hemichromis fasciatus</i> | | T | 6.5 | 11.1 | 0.0456 | 2.605 | 33 | 0.947 | Adadama Lake | this study |
| <i>Hemichromis fasciatus</i> | | T | 5.0 | 12.3 | 0.0488 | 2.495 | 138 | 0.992 | Imo River | this study |
| <i>Hemichromis fasciatus</i> | | T | 5.9 | 14.8 | 0.0102 | 3.225 | 104 | 0.992 | Mfangmfang pond | this study |
| <i>Tilapia mariae</i> | | T | 4.0 | 21.0 | 0.030 | 3.218 | 46 | 0.960 | New Calabar River | Bongonyinge (1984) |
| <i>Tilapia mariae</i> | | T | 5.0 | 25.0 | 0.7412 | 2.588 | 375 | 0.929 | Ikpa River | this study |
| <i>Tilapia mariae</i> | | T | 8.0 | 32.3 | 0.0336 | 2.852 | 51 | 0.992 | Iba-Oku stream | this study |
| <i>Tilapia mariae</i> | | T | 6.2 | 11.7 | 0.0145 | 3.169 | 211 | 0.989 | Qua Iboe River | this study |
| <i>Tilapia zillii</i> | | T | 7.0 | 15.0 | 0.0115 | 3.210 | 11 | - | New Calabar River | Bongonyinge (1984) |
| Eleotridae | | | | | | | | | | |
| <i>Bostrychus africanus</i> | | T | 4.1 | 13.0 | 0.0157 | 2.890 | 37 | 0.995 | Cross River | this study |
| Anabantidae | | | | | | | | | | |
| <i>Ctenopoma kingsleyae</i> | | T | 10.3 | 18.3 | 0.0083 | 3.364 | 65 | 0.989 | Imo River | this study |
| <i>Ctenopoma kingsleyae</i> | | T | 10.0 | 16.7 | 0.0312 | 2.840 | 146 | 0.946 | Ikpa River | this study |
| Phractolaemidae | | | | | | | | | | |
| <i>Phractolaemus ansorgei</i> | | T | 4.5 | 23.6 | 0.0061 | 3.170 | 80 | 0.972 | Imo River | this study |




Fig. 1. Distribution of the parameters of length-weight relationships in 73 populations of freshwater fishes of Nigeria. A: distribution of a values; B: distribution of b values.

Of the population studied, 50 were from lotic systems and 18 from lentic waters. Habitat-types were unspecified for five populations (see Olatunde 1985, 1989).

The results are summarized by population in Table 1. Since most of the LW data were based on relatively large samples, the resultant parameters may be considered reasonably reliable and representative. All correlations were highly significant ($P < 0.05$) with coefficients of determination ranging from 69 to 100%.

Interpopulational variability in the values of " a " was highly heterogeneous (C.V. = 282%) and ranged from $a_{\min} = 4 \times 10^{-5}$ in *Papycrocranus afer* to $a_{\max} = 7.412 \times 10^{-1}$ in *Tilapia mariae*. Conversely, interpopulational variability in " b " was low (CV = 10.8%), ranging from $b_{\min} = 2.158$ in *Nannaethips unitaeniatus* to $b_{\max} = 3.376$ in *Hepsetus odoe*. These values fall well within the limits reported by Carlander (1969, $b = 2.5-3.5$), Royce (1972, $\bar{b} = 2.0-3.5$) and Lagler et al. (1977, $b = 2.5-4.0$) for most fishes (Table 1, Fig. 1).

The mean exponent ($b = 2.911$; s.d. = 0.313) is significantly < 3 ($t = 2.405$, $df = 72$, $P < 0.02$). Torres (1991) also reported a value of $\bar{b} < 3$ in a multispecies study of LWRs. As an "assemblage", the inland water fishes of Nigeria therefore exhibit a negative allometric LWR, i.e., they tend to become thinner as they grow larger.

The exponents of fish populations inhabiting lotic ($\bar{b} = 2.911$) and lentic ($\bar{b} = 2.873$) systems were not significantly different ($t = 0.478$, $df = 66$, $P > 0.05$), thus suggesting that these broad categories of aquatic ecosystems were not different in terms of their impacts on the shape of their fish populations.

Acknowledgements

I am grateful to Mr. Okon I. Okon for field and laboratory assistance. I also thank Engr. Darling Peters for providing research fund and laboratory equipment.

References

- Beyer, J.E. 1987. On length-weight relationships. Part I: Computing the mean weight of the fish of a given length class. *Fishbyte* 5(1):11-13.
- Bolger, T. and P.L. Cannolly. 1989. The selection of suitable indices for the measurement and analysis of fish condition. *J. Fish Biol.* 34:171-182.
- Bangonyinge, C. 1984. Some observations on aspects of the biology of *Tilapia mariae* Boulenger and culture of tilapias in freshwater pond. African Regional Aquaculture Centre, Aluu, Port Harcourt, Nigeria. M. Tech. Thesis.
- Carlander, K. 1969. Handbook of freshwater fishery biology. Vol. 1. Iowa State University Press, Ames, Iowa.
- Ikusemiju, K. 1976. Distribution, reproduction and growth of the catfish, *Chrysichthys walkeri* (Günther) in the Lekki Lagoon, Nigeria. *J. Fish Biol.* 8:453-458.
- Inyang, N.M. and G.N. Anozie. 1987. Some aspects of the biology of *Epiplatys sexfasciatus* (Pisces: Cyprinodontidae) in Adada River, southeastern Nigeria. *Nigeria J. Appl. Fish. Hydrobiol.* 2:59-63.
- King, R.P. 1989. Distribution, abundance, size and feeding habits of *Brienomyrus brachyistius* (Gill, 1862) (Teleostei, Mormyridae) in a Nigerian rainforest stream. *Cybium* 13(1):23-36.
- Lagler, K.F., J.E. Bardach, R.R. Miller and D.R.M. Passino. 1977. *Ichthyology*. 2nd ed. John Wiley and Sons, New York. 506 p.
- Lévêque, C., D. Paugy and G.G. Teugels, Editors. 1992. Faune des poissons d'eaux douces et saumâtres de l'Afrique de l'Ouest (Tome 2). ORSTOM et MRAC Tervuren and Paris, 902 p.
- Nwadiaro, C.S. and P.U. Okorie. 1985. Biometric characteristics, length-weight relationship and condition factors in *Chrysichthys filamentosus* (Pisces: Bagridae) from Oguta Lake, Nigeria. *Biol. Afric.* 2(1):48-57.
- Olatunde, A.A. 1979. The biology of the family Schilbeidae (Osteichthyes: Siluriformes) in Lake Kainji, Nigeria, p. 393-412. In Proceedings of the International Conference on Kainji Lake and River Basins Development in Africa. Kainji Lake Research Institute, New Bussa, Nigeria.
- Olatunde, A.A. 1985. Length-weight relationship and the diets of *Clarias lazera* (Cuvier and Valenciennes), Family Clariidae (Osteichthyes: Siluriformes) in Zaria, Nigeria, p. 183-192. In Proceedings of the Annual Conference of the Fisheries Society of Nigeria (FISON), Kainji Lake Research Institute, New Bussa, Nigeria.
- Olatunde, A.A. 1989. Some aspects of the biology of *Synodontis schall* (Bloch & Schneider) in Zaria, Nigeria. *J. Aquat. Sci.* 4:49-54.
- Oni, S.K., J.Y. Olayemi and J.D. Adegboye. 1983. Comparative physiology of three ecologically distinct freshwater fishes, *Alestes nurse* Rüppell, *Synodontis schall* Bloch and Schneider and *Tilapia zillii* Gervais. *J. Fish Biol.* 22:105-109.
- Royce, W.F. 1972. Introduction to the fishery sciences. Academic Press, New York and London. 351 p.
- Sagua, V.O. 1986. Studies on the biology of *Gymnarchus niloticus* in Lake Chad: age determination and growth; meristic and morphometric characters, p. 179-190. In Proceedings of the Annual Conference of the Fisheries Society of Nigeria (FISON), Kainji Lake Research Institute, New Bussa, Nigeria.
- Torres, F. Jr. 1991. Tabular data on marine fishes from southern Africa, Part I: length-weight relationships. *Fishbyte* 9(1):50-53.

R.P. King is from the Department of Zoology, University of Uyo, P.M.B. 1017, Uyo, Akwa Ibom State, Nigeria.